

**Host-virus interaction: the antiviral defense function of small interfering RNAs can be enhanced by host microRNA-7 in vitro**

Xiaoying Zhang<sup>1,\*</sup>, Dongyun Liu<sup>2,\*</sup>, Sheng Zhang<sup>1</sup>, Xiujian Wei<sup>1</sup>, Jie Song<sup>1</sup>, Yupei Zhang<sup>1</sup>, Min Jin<sup>3</sup>, Zhiqiang Shen<sup>3</sup>, Xinwei Wang<sup>3</sup>, Zhichun Feng<sup>1, †</sup>, Junwen Li<sup>3, †</sup>

\* These authors contributed equally to the manuscript.

<sup>1</sup>Stem Cell Center, BaYi Children's Hospital of The General Military Hospital of Beijing PLA, 5 Nanmencang Road, Dongcheng District, Beijing, 100700, P.R.China.

<sup>2</sup>Departments of Neonatal Intensive Care Unit, The Affiliated Hospital of Qingdao University. 16 Jiangsu Road, Qingdao, 266003, P.R. China.

<sup>3</sup>Department of Environment and Health, Institute of Health and Environmental medicine, Key Laboratory of Risk Assessment and Control for Environment & Food Safety, 1 Dali Road, Heping District, Tianjin, 300050, P.R. China.

**†Corresponding author:** Prof. Zhichun Feng and Prof. Junwen Li. Prof. Zhichun Feng: 5 Nanmencang Road, Dongcheng District, Beijing, 100700, P.R.China. Phone: (86)10-66721786; Fax: (86)10-64063099; E-mail: [fengzhichun81@sina.com](mailto:fengzhichun81@sina.com). Prof. Junwen Li: 1 Dali Road, Heping District, Tianjin, 300050, P.R. China. Phone: (86)22-84655345; Fax: (86) 22-84655345; E-mail: [junwen9999@hotmail.com](mailto:junwen9999@hotmail.com)

**Supplementary Table 1. Oligonucleotide Sequences for siRNA and target genes.**


	Name	Sequence
siRNA		
5'UTR-53nt	Sense	5'-acuccgguauugcgguacccuuguatt-3'
	Antisense	5'-uacaagguaccgcaauaccggagutt-3'
5'UTR-100nt	Sense	5'-guaacuuagacgcacaaaaccaagutt-3'
	Antisense	5'-acuugguuuuugugcgucuaaguuactt-3'
5'UTR-124nt	Sense	5'-guucaauagaaggggguacaaaccatt-3'
	Antisense	5'-ugguuuguacccccuucuaauugaactt-3'
5'UTR-127nt	Sense	5'-gacggauccguuauccgcuauguatt-3'
	Antisense	5'-uacauaagcggauaacggauccguactt-3'
5'UTR-303nt	Sense	5'-gaguguagcuuaggcugaugagucutt-3'
	Antisense	5'-agacucaucagccuaagcuacacuctt-3'
5'UTR-334nt	Sense	5'-cccucaccggugacggugguccagggtt-3'
	Antisense	5'-ccuggaccaccgucaccggugagggtt-3'
5'UTR-406nt	Sense	5'-gaacaaggugugaagagccuauugatt-3'
	Antisense	5'-ucaauaggcucuucacaccuuguuctt-3'
5'UTR-472nt	Sense	5'-aaccucggagcagguggucacaaactt-3'
	Antisense	5'-guuugugaccaccugcuccgagguutt-3'
5'UTR-482nt	Sense	5'-cagguggucacaaaccagugauuggtt-3'
	Antisense	5'-ccaauacugguuugugaccaccugtt-3'
5'UTR-529nt	Sense	5'-uggcgaaccgacuacuugggugutt-3'
	Antisense	5'-acacccaaaguagucgguuccgccatt-3'
5'UTR-537nt	Sense	5'-ccgacuacuuggguguccguguuutt-3'
	Antisense	5'-aaacacggacacccaaaguagucggtt-3'
5'UTR-617nt	Sense	5'-cgaauggauuggccaucggugaatt-3'
	Antisense	5'-uucaccggauggccaauccaauucgtt-3'
Ago2	Sense	5'- ggaaaugaugcugaauauuu-3'
	Antisense	5'-auauucagcauauuuuccuu-3'
Drosha	Sense	5'-gaguaggcuucgugacuauuu-3'
	Antisense	5'-uaagucacgaagccuacucuu-3'
Negative Control	Sense	5'-uucuccgaacgugucacgutt-3'
	Antisense	5'-acgugacacguucggagaatt-3'
miRNA		
miR-7	Sense	5'-uggaagacuagugauuuuguugu-3'
	Antisense	5'-aacaauucacuagucuucauu-3'
miR-NC	Sense	5'-uucuccgaacgugucacgutt-3'
	Antisense	5'-acgugacacguucggagaatt-3'
miR-7 AS		5'-acaacaaaucacuagucuuca-3'
NC AS		5'-uuguacuacacaaaaguacug-3'
Primers		
miR-7	Stem-loop	5'-gtcgtatccagtgcgtgctgaggagtcggcaattgactgga tacgacacaaca-3'

---


	Sense	5'-cggtggaagactagtgatt-3'
	Antisense	5'-tgcgtgtcgtggagtc-3'
u6	Sense	5'-gcttcggcagcacatatactaaaat-3'
	Antisense	5'-cgcttcacgaattgcgtgcat-3'
PV	Sense	5'- ggctgctatggtgacaatcacag-3'
	Antisense	5'- gtgggtaattaatggtagaaccacc-3'
GAPDH (VERO)	Sense	5'-cgggaaactgtggcgtgat-3'
	Antisense	5'-gggtgctcgtgtgaagtcg-3'
GAPDH (A549)	Sense	5'-gaccactttgtcaagctcattcc-3'
	Antisense	5'-gtgagggtctctctctccttctgt-3'

---

# Supplementary Fig 1



## Supplementary Fig 2



### Supplementary Figure legends

**Supplementary Fig 1 Confocal microscopy analysis of efficient transfection of all the FAM-labeled siRNAs.** A549 cells were transfected with FAM-labeled siRNA-NC or FAM-labeled siRNAs, and displayed similar green fluorescent signal. Bars represent 20  $\mu$ m.

### Supplementary Fig 2 G1 arrested cells showed reduced PV infection.

(a) A549 cells are arrested in G1 by siRNA directed against cdk2 (gene ID: 1017, target sequence: AGGTGGTGGCGCTTAAGAAAA) and cyclin A (gene ID: 890, target sequence: GCCAGCTGTCAGGATAATAAA) before PV infection as analyzed by flow cytometry. (b) The relative expression of PV in

different siRNA transfected cells determined by real-time PCR. PV titer:  $10^{-8}$

TCID<sub>50</sub>/100 μL.